

**MESTRADO INTEGRADO EM MEDICINA**

2016/2017

Helena Margarida Cruz Gens  
CCR na mulher grávida  
CRC in pregnant women

março, 2017

FMUP

Helena Margarida Cruz Gens  
CCR na mulher grávida  
CRC in pregnant women

**Mestrado Integrado em Medicina**

**Área: Cirurgia Geral**  
**Tipologia: Monografia**

**Trabalho efetuado sob a Orientação de:**  
**Mestre Laura Elisabete Ribeiro**  
**Barbosa**

**Trabalho organizado de acordo com as normas da revista:**  
**Journal of Coloproctology**

março, 2017

**FMUP**

Eu, Helena Margarida Cruz Gomes, abaixo assinado, nº mecanográfico 201104854, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 13 / 03 / 2017

Assinatura conforme cartão de identificação:

Helena Margarida Cruz Gomes

NOME

Helena Mangualda Cruz Gens

NÚMERO DE ESTUDANTE

201104854

E-MAIL

helenagem93@gmail.com

DESIGNAÇÃO DA ÁREA DO PROJECTO

Cirurgia Geral

TÍTULO DISSERTAÇÃO/MONOGRAFIA (riscar o que não interessa)

CCR na mulher grávida / CRC in pregnant women

ORIENTADOR

Luísa Elisabete Ribeiro Barbosa

COORDINADOR (se aplicável)

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTES TRABALHOS APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input checked="" type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTES TRABALHOS (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTES TRABALHOS.	<input type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 22/03/17

Assinatura conforme cartão de identificação: Helena Mangualda Cruz Gens

# CCR na mulher grávida

## CRC in pregnant women

Helena Margarida Cruz Gens<sup>1</sup>

Laura Elisabete Ribeiro Barbosa<sup>2</sup>

<sup>1</sup>Universidade do Porto, Faculdade de Medicina, Porto, Portugal

<sup>2</sup>Universidade do Porto, Faculdade de Medicina, Porto, Portugal, Centro Hospitalar São João, Serviço de Cirurgia Geral, Porto, Portugal

Helena Margarida Cruz Gens

Rua Ernesto Santos Ribeiro, 196

4470-176 Maia

[helenagens93@gmail.com](mailto:helenagens93@gmail.com)

## Abstract

### **Background**

Colorectal cancer in pregnancy is a rare pathology with limited high-grade evidence available for guidance. The diagnosis of CRC in pregnant women is usually delayed, and once diagnosis is made, challenges exist as treatment options may be limited.

### **Objective**

The study aims to highlight the importance of early investigation of symptomatic patients during pregnancy, as well as to update treatment and prognosis in CRC.

### **Methods**

A literature search in Pubmed database, including articles from 2006 to 2016 and cross-research articles with the initial research.

### **Results**

Pregnancy can limit and contraindicate the utilization of standard diagnostic and therapeutic tools, which in particular can hamper the liberal use of colonoscopy and CT. Physical evaluation and abdominal US are first recommended; besides, MRI or CT may be used, only in indicated cases.

Surgery is the main stay of treatment but radiotherapy and chemotherapy have significant role in posterior management of tumour.

### **Discussion**

As abdominal symptoms are common in pregnancy and colorectal cancer can simulate them, the differential diagnosis between these two pathologies is crucial, since early interventions can be curative.

After diagnosis, the follow-up of pregnant women should be individualized, depending on several factors. However, since cancer treatment is currently possible in pregnancy, the patient's survival probability should not be decreased due to late diagnosis.

### **Conclusions**

Many studies are needed in order to achieve development in CRC pathogenesis during pregnancy as well as in treatment outcomes.

The potential curative treatment of the disease should be the main aim of treatment when considering CRC in pregnancy. However, it is crucial to adapt the treatment to each patient, taking into account conscious decision on pregnancy further management.

## Resumo

### Introdução

O cancro colorretal na gravidez é uma patologia rara, com limitada evidência científica para orientação terapêutica. O diagnóstico de cancro colorretal em mulheres grávidas é tardio e, quando o diagnóstico é feito, as opções de tratamento podem ser limitadas.

### Objetivo

O objetivo deste estudo é ressaltar a importância da investigação precoce de pacientes sintomáticas durante a gravidez, assim como avaliar os atuais métodos de tratamento e prognóstico no CCR.

### Métodos

A pesquisa bibliográfica foi realizada na base de dados PubMed, incluindo artigos a partir de 2006 até 2016, assim como artigos de pesquisa cruzada com os artigos iniciais.

### Resultados

A gravidez pode limitar e contra-indicar a utilização de ferramentas de diagnóstico e terapêuticas convencionais, assim como dificultar o uso indiscriminado de colonoscopia e tomografia computadorizada. A avaliação física e a ecografia abdominal são a primeira linha para diagnóstico. No entanto, em casos selecionados, a ressonância magnética ou a tomografia computadorizada também podem ser usadas.

A cirurgia é o *gold-standard*, mas a radioterapia e a quimioterapia assumem um papel cada vez mais relevante no tratamento multidisciplinar destes tumores.

### Discussão

Como os sintomas abdominais são comuns na gravidez e o cancro colorretal pode simulá-los, o diagnóstico diferencial entre estas duas patologias é crucial, já que intervenções precoces podem ser curativas.

Após diagnóstico, o seguimento das grávidas deve ser individualizado, dependendo de vários fatores. Porém, já que atualmente o tratamento do cancro é possível na gravidez, a probabilidade de sobrevivência da paciente não deve ser diminuída devido ao diagnóstico tardio.

### Conclusões

Outros estudos são ainda necessários para saber mais acerca da patogénese do cancro colorretal na gravidez, assim como os resultados após tratamento.

O potencial objetivo é o tratamento oncológico do cancro colorretal. No entanto, é crucial adaptar o tratamento a cada paciente, tendo em conta a sua decisão consciente acerca da continuação da gravidez.

Key-words

Cancro do cólon; Cancro do reto; Gravidez; Diagnóstico do cancro; Tratamento do cancro.

Palavras chave

Colon cancer; Rectal cancer; Pregnancy; Cancer diagnosis; Cancer treatment.



## Introduction

### Background

Colorectal cancer (CRC) in pregnancy is a condition that represents a distinct entity from CRC in the general population, as it is a rare pathology with limited high-grade evidence available for guidance. In fact, the literature on this subject is scant with fewer than 300 cases reported(1), which means there is limited experience on the management of CRC diagnosed during pregnancy.

The diagnosis of CRC in pregnant women is usually delayed, because there is a significant overlap in signs and symptoms between a colorectal malignancy and normal pregnancy, impeding proper diagnosis(2–4). Once diagnosis is made, challenges exist as treatment options may be limited(2). That is why management requires judiciously individualized strategies after thorough patient counselling to deal with consequent emotional and physical stress, in order to ensure adequate psychological support and realistic expectations(2,5).

The primary objective of this review was to highlight the importance of early investigation of symptomatic patients during pregnancy, as well as the role of chemotherapeutic, radiological and surgical interventions. Secondary objectives include a research about epidemiology of CRC during pregnancy, its implications and prognostic features.

### Methods

The literature survey was conducted in Pubmed database. The words "colon pregnancy cancer" and "rectum pregnancy cancer" were used. Only the articles published from 2006 to 2016 were considered. After reading the title and abstract, and subject to an availability of the article, 31 articles were obtained in Pubmed. Articles obtained by cross-searching with the articles of the initial research with relevant information were also added.

## Results

### Epidemiology

CRC is the third most common type of cancer in women(4) with its highest incidence occurring in patients aged 50 years old(6); women older than 40 years tend to be 11 times more likely to develop CRC than those younger than 30 years of age(7,8).

While the overall incidence of CRC is steady or falling, some studies report an increased incidence of CRC in younger patients (< 40 years)(1), which means that 3% of patients with this cancer are younger than 40 years old(6). Nevertheless, for this age group, studies report that overall survival of CRC for women has improved substantially, with 5-year overall survival now approaching 80%(9).

Because of the increase in the incidence of CRC in younger patients, it may occur during the reproductive age, interfering with pregnancy(1). Cruveilhier reported the first case of rectal carcinoma in pregnancy in 1842, but now it is the seventh most common type of cancer in pregnancy(5,6). However, its incidence rate is controversial; some studies report an incidence of 0,002%(3,4,6–8,10,11) while others consider 0.07% to 0,1%(12–14), which means about 1 case per 13.000 gestations(3,5,15). The mean age of women with CRC during pregnancy has been reported as 31 years of age(4–6,16),

with range of 16-48 years(12)

Considering racial predisposition, hispanic women were slightly less likely to develop CRC, in comparison with non-hispanic white women(8).

Most of CRC in pregnancy is an aggressive mucinous subtype(8), which have poorer prognosis(17), but primary signet-cell carcinoma (SRCC) of the colon and rectum also represents a form of adenocarcinoma of the large intestine. Although its rare incidence of about less than 0,1% of cases of CRC in pregnancy, patients with SRCC are younger(14).

### Etiology and pathogenesis of CRC in pregnancy

CRC is one of the three most common types of cancer in women and can present in women of childbearing age, especially if there is a genetic predisposition(12).

In fact, environmental factors play a dominant role in the etiology of most CRC but inherited genetic factors are also significant in between 15% and 30% of cases. In about 5% of all cases, CRC is associated with a highly penetrant dominant or recessive inherited syndrome(18). When considering familial clustering of CRC and endometrial cancers, it is important to consider Lynch's syndrome (hereditary non-polyposis colorectal cancer (HNPCC)), as a cause of cancer in pregnant women(19). It is an autosomal dominant inherited genetic disease(20), and thus multiple generations can develop CRC at an early age (mean, 45 years). Lynch syndrome is likely if a family history meets the Modified Amsterdam Criteria or revised Bethesda guidelines(11)

Familial adenomatous polyposis (FAP) is another inherited syndrome, responsible for <1% of all CRC cases. FAP is transmitted as an autosomal dominant trait, and is caused by truncating mutations in the (APC) adenomatosis polyposis coli gene. Recently, the **MUTYH** (mutY homologue (**Escherichia coli**)) gene has been identified as a further polyposis gene, displaying an autosomal recessive pattern of inheritance(18).

In what concerns to environmental factors, delayed childbearing and increased maternal age may lead to an increased incidence of CRC complicating pregnancy(6,21).

Some investigators demonstrated that 20 to 54% of colon cancers have estrogen receptors (ERs), whereas others have demonstrated progesterone receptors (PgRs), which may be stimulated by the estrogen and progesterone produced during pregnancy. The role of these hormones in the etiology and progression of CRC are limited and conflicting(12). In fact, CRC pathogenesis and its relation to pregnancy is not well understood(8,12,17), and studies show that parity is not positively neither negatively associated with CRC(8).

When discovered during pregnancy, two-thirds of CRC in pregnant women tend to involve the rectum and sigmoid colon, unlike the general population where two-thirds arise from the extra pelvic colon(4,12–14,17). In fact, it was reported that about 85% of CRC in pregnancy are below the peritoneal reflection(1,5,12,16).

### Prevention and screening

Advances in molecular basis of CRC include identifying the adenomatous polyposis coli (APC) gene, P<sup>53</sup> gene, mismatch repair genes, and loss of allelic heterozygosity.(17)

Accordingly, as familial adenomatous polyposis (PAF) is a known risk factor for CRC during pregnancy, patients with family history of HNPCC should perform genetic testing(1,5).

When a patient develop CRC at a young age, it is important to consider the possibility of a hereditary cause, so we should confirm her family medical history in what concerns to cancers. Identification of the germline mutation in a Lynch syndrome family allows their inclusion in lifesaving cancer surveillance programs, which has been proven to reduce CRC mortality(22). Therefore, screening tests should be performed on tumour tissues to help determine the likelihood of this condition and microsatellite instability (MSI) analysis is the first approach to identify patients with Lynch syndrome. Germline testing for mismatch repair (MMR) gene alterations should be performed(11), as an autosomal dominant MMR deficiency leading to a tumour with MSI was assumed to be the primary mechanism for Lynch syndrome(23). In fact, germline mutations in the genes MLH1, MLH2, MSH6 and PMS2 can lead to the development of the syndrome, and heterozygosity for a mutation in one of these genes can result in increased susceptibility to cancer(11).

If Lynch syndrome had been suspected early diagnosis is essential. The American Cancer Society guidelines recommend colonoscopy beginning at an earlier age for high-risk individuals(11,24). Periodic examination by colonoscopy leads to the detection of CRC at an earlier stage, to a 63% reduction of the risk of CRC and to a significant reduction of the mortality associated(25). Annual colonoscopy programs performed at the age of 25 years in patients with families that have at least 3 relatives with a history of CRC or other HNPCC-related tumours and in families with identified MMR defect(18). It is recommended a 3-year gap between colonoscopies because this time interval has proven effective for the detection of this condition(11).

When pregnant women with SRCC are analysed for microsatellite instability, studies confirm they represent about 30% of tumours. Moreover, mutations of K-ras and P<sup>53</sup> gene have been reported in SRCC. Furthermore, replications of DNA are also suggested to be at least partly involved in carcinogenesis of SRCC(14).

Note down that genetic testing is only recommended for those with a family history suggestive of Lynch syndrome or other hereditary pattern, as most CRC are sporadic(11).

### Clinical features and Diagnosis

Diagnosis of CRC during pregnancy is challenging secondary to the difficulty in distinguishing pain of gynaecologic and GI origin(16,26), and the overlap in signs and symptoms of cancer and pregnancy(1–3,5,13,16). Consequently, usually there is a delay in diagnosis of CRC during pregnancy(4–7,10,12,26). Those signs and symptoms include nausea, vomiting, abdominal pain, weight loss, anemia, abdominal mass, rectal bleeding and altered bowel habits such as constipation.

Physiologic adaptations in pregnancy may also alter clinical presentations(12) so, high degree of suspicion if clinical features are suggestive of GI obstruction.(17)

However, sometimes there is no suspicion of CRC because its symptoms are absent(3) and foetal movements are normal(17). In fact, several studies reported that CRC can develop during pregnancy without presenting symptoms(6,7,27).

When considering SRCC, symptoms usually develop later, leading to the non-

detection of cancers limited to mucosa and sub-mucosa(14).

Palpable abdominal mass is a uncommon finding(17) but a common problem can be excluded with a careful anorectal exam(1), yielding 67-84% accuracy in staging rectal carcinomas(14).

Intestinal obstructions are extremely rare in pregnant women<sup>(2)</sup> but when present, exploratory laparotomy must be commenced(28), as prompt operative intervention maximizes outcome for both foetus and mother(3).

Haemorrhoids or anal fissure, common causes of rectal bleeding, should be evaluated, and rectal examination should be performed when a patient presents with a complaint of pain and/or rectal bleeding(6,13). Persistent anorectal bleeding or rectal passage of tissue at the time of delivery is an ominous sign of CRC cancer and should be investigated(12).

If a patient starts to lose weight while pregnant, she should be evaluated for maternal and foetal etiologies. Persistent nausea or vomiting, specially in the third trimester, should also be evaluated further(6).

Above all, it is crucial to consider complaints that are specific, severe or perseverate(3,4).

The diagnosis of CRC in a non-pregnant patient entails the tumour marker serum carcinoembryonic antigen (CEA), toraco-abdomino-pelvic imaging, and endoscopy with biopsy.

- CEA

Evaluation of abnormal laboratory values is important to optimize patient diagnosis. CEA levels have been used during pregnancy for the diagnosis, monitoring and prognosis of CRC. Unfortunately, CEA levels tend to be normal or slightly elevated during pregnancy(14,16,17), and are not considered a useful screening tool due to their low sensivity and specificity(12).

- Colonoscopy

Colonoscopy is the gold standard to confirm diagnosis as it provides direct visualization, accurate determination of location, and the opportunity to obtain tumour tissue for pathological diagnosis of CRC(4,14). The American Society of Gastrointestinal Endoscopy indicate that an endoscopic intervention is safer than radiologically guided or surgical operations(1,5). However, pregnancy is a relative contraindication as this procedure may complicate pregnancy(4,12–14), due to the risk of foetal exposure to potential teratogenic medications, uteroplacental insufficiency with maternal hypoxia or hypotention, and the risk for placental abruption with the mechanical pressure to the uterus(1,2,5,12). Accordingly, it is advised that strong indication should be present to proceed with an endoscopy, it should be delayed until second trimester where possible(1,5), the procedure time must be minimized and the lowest possible dose of sedative medications should be used(2).

With informed patient approval, the procedure may be performed with possible reduction in risk with the use of meperidine, because of its safer foetal profile, maternal oxygen administration, and gentle abdominal compression(12,14).

Because most cases of CRC during pregnancy are rectal carcinomas, if the lesion is confined to the distal colon, a flexible rectosigmoidoscopy (preferably without sedation) may be performed, as an alternative to colonoscopy(1,5,12,13). These one could be delayed in order to look for synchronous lesions.

## Staging

In the literature, most CRC are usually detected in its advanced stages, secondary to late diagnosis(3), during second or third trimester of pregnancy(6,12,13). A study that reviewed 41 patients with CRC during pregnancy observed that all patients had stage II or greater disease(12), while other refers that 60% are already diagnosed at stage III or IV(13). It corresponds to Dukes stage B or higher at presentation(4).

Staging is critical to ensure one is not dealing with advanced stage IV disease, as local management of colorectal malignancy would change considerably(2).

- CT scan

Imaging evaluation during pregnancy is difficult since toraco-abdomino-pelvic CT scan, is relatively contraindicated in pregnancy. This procedure should be avoided particularly in the first trimester, secondary to the foetal risk of radiation exposure teratogenicity and carcinogenicity(2–4,12–14,17). However, it is suggested that risk of adverse foetal effects is very low at doses of radiation used for diagnostic purposes, so CT of pelvis and abdomen can be performed with minimal risk(2,28).

- Ultrasound

Ultrasound (US) evaluation is a reasonable alternative to CT. (17) It is especially useful for detection of hepatic metastases (2) (75% of sensitivity)(12,14) but, because of the gravid uterus, it has limited accuracy in detecting colon and rectal masses (3,12,13). Further diagnostics are frequently necessary when ultrasounds are negative.(3)

Transrectal ultrasound is helpful in late pregnancy to determine rectal cancer location and is more accurate in staging rectal cancer preoperatively, although it has not been proven to prolong survival.(17)

- MRI

Similarly, magnetic resonance imaging (MRI) is relatively safe in pregnancy and should be considered after US indeterminate findings (28), but is good practice to avoid non-urgent MRI, particularly in the first trimester. (1,2) Moreover, it should be performed without contrast (3) because contrasts have not been approved for the foetus (12,13), being gadolinium known as teratogenic agent. (2)

## Treatment

Treatment during pregnancy is another challenging issue; Walsh et al have proposed an algorithm to manage CRC diagnosed during pregnancy based on the gestational age of the foetus, foetal lung maturity, cancer stage, need for adjuvant chemotherapy, and if elective or emergent surgery is indicated.(1,5,6,12,14,15)

The treatment goal is to implement therapy as soon as possible for the mother, (6,13) and balance this with delivery of the foetus with optimal for neonatal outcome. (1,12,13,16)

Treatment modalities may include surgery, radiation therapy and chemotherapy, depending on the stage of the cancer.(6)

- Surgery

Surgery is the primary therapy for CRC outside of pregnancy (12,14) (8) In pregnant women with CRC it is recognized as safe and feasible (1) but its timing in a is a pivotal issue (2) so it should be considered on a case-by-case basis.

However, as the majority of CRC malignancies diagnosed in pregnancy occur below the peritoneal reflection, the level of technical difficulty associated with surgery is increased.(2)

If bowel obstruction develops during the pregnancy, a self-expanding metallic stent (SEMS) can be useful as it allows solving the acute condition, providing time to prepare the patient for surgery, thus reducing both post-surgical morbidity and mortality.(29)

Although no high-grade evidence exists regarding management of CRC in pregnancy, some clinical guidelines are reported in literature:

- If diagnosis is made in the first 20 weeks of pregnancy, treatment delay can lead to disease progression and compromise of mother's life. Therefore, the recommendation is discontinuation of the pregnancy, according mother's followed by early cancer treatment with surgical resection, as in non-pregnant patients. (2,13,14,16) However, controversia data regarding risk to the pregnancy with surgery, if imaging suggests that the tumour may be resected with clear margins, surgery might be an option. (2) In fact, low anterior or abdominoperineal resection has already been performed up to 20 weeks gestation without disturbance of the gravid uterus.(12) However, as CRC diagnosis is rarely made prior to 20 weeks of gestation, there is limited data on foetal outcome after surgical resection.(12)

- If colon cancer is diagnosed after 20 weeks gestation, surgery can be delayed until delivery, in order to save the foetus (5,13,14,16), although endangering the patient with significant risk of disease progression (3), due to the pro-angiogenic state of pregnancy.(2) The ultimate goal is to achieve foetal lung maturation (8,16); nevertheless, delivery may vary from 28 to 32 weeks gestation, based on multispecialty team decision.(2,6,12) After that, treatment of CRC should take place as in a non-pregnant patient.(2,14,15)

In both cases, the extent of resection is determined by tumour size, location, histologic grade and tumour extension into the colon wall and into adjacent tissue and organs.(14)

- Chemotherapy

The need for chemotherapy depends upon the final histology of the tumour (2), being considered in stage II with high risk of recurrence (14) and stage III when nodal involvement is present.(8)

As a large portion of patients are diagnosed in advanced stages, it is common that neoadjuvant chemotherapy before surgical resection is needed in rectal cancer. (13) Adjuvant chemotherapy has been shown to improve the survival rate by 5 to 10% for stage II or III CRC, (12) but evidence shows that in pregnant women with metastatic rectal cancer it might spare the foetus, but not cure the mothers. (13)

Moreover, pelvic radiation is not recommended during organogenesis, in the first trimester of pregnancy (5,15,16), as it is associated with lethal damage to the foetus, with embryonic or foetal death, malformation, and growth retardation. (13,15) Although some studies reported that chemotherapy should be given only after delivery, other ones suggest that chemotherapy can be administered in the second or the third trimesters with dose maternal/foetal surveillance. (1,5,15) The recommended therapeutic agent is 5-fluoruracil (5-FU), which is an inhibitor of DNA synthesis.(4) Although 5-FU is reported to be associated with low or no risk of adverse reproductive outcomes (9,10), some investigators have suggested a the possibility of spontaneous abortion (1,5) and teratogenicity associated with 5-FU (5,12,15,16)

Other new chemotherapeutic agents platinum-based like cisplatin and oxaliplatin are available but, according to the U. S. Food and Drug Administration, they are not recommended during pregnancy. (12,15,16)

The previous reasons could explain why women with pregnancy-associated CRC were less likely to undergo chemotherapy, in comparison to non-pregnant women in the same condition.(8)

It is also important to report that, considering Lynch syndrome, tumours with MSI were more responsive to adjuvant chemotherapy than tumours without MSI.(11)

- Radiotherapy

Adjuvant radiotherapy is indicated for Duke's B2 and C rectal cancers (12), T4 lesions adherent to the pelvic structure and in patients with close or positive surgical margins.(8,14) Nevertheless, radiation treatment of the pelvis is contraindicated during pregnancy (2,5,15) and is usually delayed until after delivery (12,16) as it has been implicated in sexual and gonadal dysfunction, foetal growth restriction or spontaneous miscarriage. (9)

- Biological therapies

Biological agents like bevacizumab, cetuximab or panitumumab provide relatively modest survival increase in addition to standard chemotherapy, withholding their use until delivery would not be likely to prevent curative treatment.(10)

In stage IV disease, palliative management should be performed and emphasis should be to lengthen the progression-free and overall survival in the unresectable metastatic CRC. (14)

(12)

Vaginal delivery vs cesarean

The delivery mode is controversial as cancer per se is not an indication to perform a cesarean section. (5,16) However studies show that cesarean section is more practiced in women with pregnancy associated with CRC.(8)

Outside the normal obstetrical indications for cesarean section, indications for an operative delivery in CRC patients include a tumour along the anterior rectal wall, secondary to increased risk of bleeding with vaginal birth pressure or birth canal obstruction by tumour.(12) Cesarean also can be considered if resection of the CRC at the time of delivery will be performed.(5,14,16)

There has been a recent move toward vaginal delivery for women with rectal carcinoma, even with an unripe cervix requiring a cervical ripening agent. (12)

### Complications

The delay in diagnosis lead to an increase of complications as the uterus, cervix and adnexa share the same visceral innervation as the lower ileum, sigmoid colon, and rectum. (26) About 25% of pregnant women with CRC will have ovarian metastases. (12,14,16) If tumor resection is performed during pregnancy, prophylactic ovarian removal may be deferred secondary to the possible risk for a spontaneous abortion, especially in the first trimester. Bilateral oophorectomy is performed during pregnancy, if evidence of invasion. (12,16)

The liver is the most common site for synchronous metastases (14) but there are no reports of liver resection for CRC liver metastases during pregnancy.(10,16)

Colon obstruction, perforation, and metastasis are more frequent in pregnant women with CRC than the average population, (17) possibly due to the immunosuppressive state of pregnancy.(12)

### Foetal risk

It is reported that higher rates of preterm labour and major puerperal infections are noted in women with pregnancy associated with CRC, as this malignant condition is prone to infections that may be sub-clinical before delivery. (8) This could be explained by the malignancy-related immune suppression, as CRC initiates an inflammatory reaction that consequently starts the preterm labour cascade secondarily to the close proximity to the uterus.(8)

Despite high rates of preterm delivery, some studies reported an absence of adverse foetal outcomes (5,8), while others refer that only 78% of foetus from women with CRC survived.(5) Conversely, there is agreement that the risk of foetal malignancy is rarely observed, even when the disease is in an advanced stage with widespread metastasis. (5,17) However, metastasis to the placenta was reported once in maternal colorectal malignancy.

Even though a complete evaluation of the placenta is recommended, there is no evidence to support periodic follow-up of the baby.(5)

### Prognosis

Pregnant women with CRC tend to have poor outcome, which usually includes widespread metastasis (5), mainly secondary to delayed diagnosis. (4,6,8,12,16).



Moreover, when considering SRCC, it is more likely to experience lymph node metastasis, have an aggressive clinical course and poor prognosis. (14)

Previous studies reported that patients with CCR regarding Lynch syndrome with MSI survived longer than patients with non-MSI tumours did; accordingly, the former ones showed lower mortality rates when stratified by tumour stage, including patients with stage IV cancer.(11) Hence, the detection of MSI in a CCR is a positive prognostic factor, particularly among young patients.(11,20)

Most of all, it is important to notice that women that are not surgically treated died at a rate that was 4.2 times that of women who undergo surgical treatment.(8)

When comparing pregnant to non-pregnant women with CRC, 5 year cancer survival is the same (12) Women with colon cancer died at a faster rate than those with rectal or anal cancer, even though it is reported that pregnancy was not associated with a significant difference in survival between these two groups. (5,8)

## Discussion

After this review it appears that there is still much to be clarified, especially in the treatment area. Despite a rising incidence of CRC in pregnancy due to, at least in part, more women falling pregnant at an advanced maternal age, only few studies were performed about this issue. Moreover, most of those have not been completed due to ethical reasons, so there is information evaluating new therapies.

CRC in pregnancy is rare but its incidence is increasing not only because more women are postponing pregnancy until later in life but also because there is an increased risk for this cancer in women with more than 40 years old.(5,10)

This cancer in gravidity is not well understood but, as genetic alterations contribute to the susceptibility to CRC it is important to offer genetic counselling, especially when other CRC is recognised in the family history.(8)

As abdominal symptoms are common in pregnancy and CRC can mimic them, they are overlooked. (6,13,16) Thus, differential diagnosis should include the patient's history, physical examination, laboratory data, and radiologic findings that may assist in identifying the diagnosis (12); an appropriate work-up may result in prompt evaluation, detection and appropriate interventions to treat CRC (6,12,28) as early diagnosis improves survival and treatment outcomes.(1,2,6) In the era of possible cancer treatment in pregnancy, patient's chance of survival should not be diminished by delayed diagnosis.(7)

Further management is individualized and dependent on various factors including maternal age, patient's desire for future fertility, gestational age at diagnosis, and cancer stage.(12) However, pregnancy can limit and contraindicate the utilization of standard diagnostic and therapeutic tools due to a gravid uterus and a potentially vulnerable foetus, which can hamper the liberal use of colonoscopy and CT. (1,13) Physical evaluation plays an essential role, specially the evaluation of liver size and a rectal examination to screen for masses. (6) Abdominal US is first recommended, but its use is limited due to the patient's change in body habitus (28). MRI or CT may be used if necessary, although CT is not desirable due to foetal irradiation (28). Laparoscopy,

EGD, and sigmoidoscopy can be performed during pregnancy, when strongly indicated.(12)

Most medications appear to be relatively safe to the foetus and can be used when benefits to the mother outweigh potential foetal risks. (12)

Surgery is the main stay of treatment but its indications differ if CRC diagnostic occurs in the first half or in the second half of pregnancy. Secondly, in adjuvant and neoadjuvant settings, RT and CRT have significant role in accordance to the site of tumor.(14) However, there is few data suggest association between pregnancy outcome and treatment with chemotherapeutic agents, used in treatment of CRC. (12) Therefore, it may still difficult to provide precise guidance to patients about long-term effects of this treatment. (9,16)

## Conclusion

The coexistence of malignant tumour and pregnancy is a state of simultaneous occurrence of two completely contradictory phenomena – the development of a new life and a life-threatening terminal illness. In fact, CRC is an aggressive cancer that is rarely found during pregnancy, but when it appears it is expected poor outcome, as its usually diagnosed in late stages. (6,12,13)

Because there are no absolute guidelines, it is associated with diagnostic and therapeutic challenges,(10,13) as gravidity requires further quick and adequate diagnosis. (3)

There is necessary further investigations about diagnostic and treatment modalities with reduced foetal side effects, in order to diminish its incidence and mortality rate.(1) Follow-up of the infants in later childhood and adolescence as the central nervous system continue to develop, with additional reporting of cases, is needed to establish the safety of chemotherapeutic treatment of CRC during pregnancy.(16)

In fact, treatment during pregnancy varies widely and poses significant legal, ethical, religious, emotional and scientific challenges; therapy should be individualized and defined by a multidisciplinary team (12) that considers not only through patient counselling, but also the best management for both the patient and her foetus. (2) All the professionals who look after such a special patient should inform her considering the most current and reliable knowledge, providing her a multidisciplinary care, and understanding the complexity of coexistence of cancer and pregnancy. The patient should have the opportunity to decide about the fate of her pregnancy and it should not be affected by the moral beliefs of the doctor; the final decision concerns only to the patient(1,21,30,31)

Above all, the treatment strategy for CRC should be no different for pregnant and non-pregnant patients in terms of the aim, which is potential curative treatment of the disease, but should always consider the patient's conscious decision on pregnancy further management.

### Conflict of interest

The authors have no commercial or other associations that might pose a conflict of interest in connection with the manuscript.

## References

1. Aytac E, Ozuner G, Isik O, Gorgun E, Stocchi L. Management of colorectal neoplasia during pregnancy and in the postpartum period. *World J Gastrointest Oncol*. 15 de Julho de 2016;8(7):550–4.
2. Bukhari Y, Hogan NM, Pomeroy M, O’Leary M, Joyce MR. Surgical management of rectal cancer in pregnancy. *Int J Colorectal Dis*. 22 de Julho de 2012;28(6):883–4.
3. Ossendorp RR, Silvis R, van der Bij GJ. Advanced colorectal cancer resulting in acute bowel obstruction during pregnancy; a case report. *Ann Med Surg*. Junho de 2016;8:18–20.
4. Kraljević M, Hoffmann H, Knipprath A, von Holzen U. Obstructing adenocarcinoma of the descending colon in a 31-year-old pregnant woman. *Int J Surg Case Rep*. 16 de Outubro de 2014;5(12):958–60.
5. Yaghoobi M, Koren G, Nulman I. Challenges to diagnosing colorectal cancer during pregnancy. *Can Fam Physician*. Setembro de 2009;55(9):881–5.
6. Khangura RK, Khangura CK, Desai A, Goyert G, Sangha R. Metastatic Colorectal Cancer Resembling Severe Preeclampsia in Pregnancy. *Case Rep Obstet Gynecol* [Internet]. 2015 [citado 23 de Setembro de 2016];2015. Disponível em: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4685096/>
7. Gabriel I, Olejek A, Drozdowska B. Colon cancer in pregnancy - a difficult diagnosis. *Eur J Obstet Gynecol Reprod Biol*. Agosto de 2016;203:340–1.
8. Dahling MT, Xing G, Cress R, Daniels B, Smith LH. Pregnancy-associated colon and rectal cancer: perinatal and cancer outcomes. *J Matern-Fetal Neonatal Med Off J Eur Assoc Perinat Med Fed Asia Ocean Perinat Soc Int Soc Perinat Obstet*. Março de 2009;22(3):204–11.
9. Haggar F, Pereira G, Preen D, Woods J, Martel G, Boushey R, et al. Maternal and neonatal outcomes in pregnancies following colorectal cancer. *Surg Endosc*. Julho de 2013;27(7):2327–36.
10. Robson DE, Lewin J, Cheng AW, O’Rourke NA, Cavallucci DJ. Synchronous colorectal liver metastases in pregnancy and post-partum: sCRLMs in pregnancy and post-partum. *ANZ J Surg*. Julho de 2015;n/a-n/a.
11. Okuda T, Ishii H, Yamashita S, Ijichi S, Matsuo S, Okimura H, et al. Rectal Cancer Diagnosed after Cesarean Section in Which High Microsatellite Instability Indicated the Presence of Lynch Syndrome. *Case Rep Obstet Gynecol* [Internet]. 2015 [citado 19 de Outubro de 2016];2015. Disponível em: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4439484/>
12. Longo SA, Moore RC, Canzonieri BJ, Robichaux A. Gastrointestinal Conditions during Pregnancy. *Clin Colon Rectal Surg*. Junho de 2010;23(2):80–9.

13. Toosi M, Moaddabshoar L, Malek-Hosseini SA, Sasani MR, Mokhtari M, Mohammadianpanah M. Rectal cancer in pregnancy: A diagnostic and therapeutic challenge. *J Egypt Natl Cancer Inst.* Setembro de 2014;26(3):175–9.
14. Dhull AK, Gogia P, Atri R, Dhankhar R, Kaushal V, Singh S, et al. Exploring signet-ring cells in pregnant female. *J Gastrointest Oncol.* Abril de 2015;6(2):E10–5.
15. Basta P, Bak A, Roszkowski K. Cancer treatment in pregnant women. *Contemp Oncol.* 2015;19(5):354–60.
16. Makoshi Z, Perrott C, Al-Khatani K, Al-Mohaisen F. Chemotherapeutic treatment of colorectal cancer in pregnancy: case report. *J Med Case Reports* [Internet]. 13 de Junho de 2015 [citado 22 de Setembro de 2016];9. Disponível em: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4469397/>
17. Mechery J, Ikhenia SE. Cancer of the descending colon during pregnancy. *J Obstet Gynaecol.* Janeiro de 2007;27(3):311–2.
18. Vasen HFA, Moslein G, Alonso A, Bernstein I, Bertario L, Blanco I, et al. Guidelines for the clinical management of Lynch syndrome (hereditary non-polyposis cancer). *J Med Genet.* 27 de Fevereiro de 2007;44(6):353–62.
19. Lynch HT, Lynch JF, Attard TA. Diagnosis and management of hereditary colorectal cancer syndromes: Lynch syndrome as a model. *CMAJ Can Med Assoc J.* 1 de Setembro de 2009;181(5):273–80.
20. Boland CR, Goel A. Microsatellite Instability in Colorectal Cancer. *Gastroenterology.* Junho de 2010;138(6):2073–2087.e3.
21. Skrzypczyk-Ostaszewicz A, Rubach M. Gynaecological cancers coexisting with pregnancy – a literature review. *Współczesna Onkol.* 2016;3:193–8.
22. Akoum R, Ghaoui A, Brihi E, Ghabash M, Hajjar N. Early-onset breast cancer in a Lebanese family with Lynch syndrome due to MSH2 gene mutation. *Hered Cancer Clin Pract* [Internet]. Dezembro de 2009 [citado 22 de Outubro de 2016];7(1). Disponível em: <http://hccpjournal.biomedcentral.com/articles/10.1186/1897-4287-7-10>
23. Bansidhar BJ, Silinsky J. History and Pathogenesis of Lynch Syndrome. *Clin Colon Rectal Surg.* Junho de 2012;25(2):63–6.
24. Smith RA, Cokkinides V, Brooks D, Saslow D, Shah M, Brawley OW. Cancer screening in the United States, 2011: A Review of Current American Cancer Society Guidelines and Issues in Cancer Screening. *CA Cancer J Clin.* Janeiro de 2011;61(1):8–30.
25. Engel C, Rahner N, Schulmann K, Holinski-Feder E, Goecke TO, Schackert HK, et al. Efficacy of Annual Colonoscopic Surveillance in Individuals With Hereditary Nonpolyposis Colorectal Cancer. *Clin Gastroenterol Hepatol.* Fevereiro de 2010;8(2):174–82.

26. Augustin G, Majerovic M. Non-obstetrical acute abdomen during pregnancy. *Eur J Obstet Gynecol Reprod Biol.* Março de 2007;131(1):4–12.
27. Castresana D, Wong M, Kaza A. Rectal Mass in a Pregnant Patient: How Come? *Dig Dis Sci.* Dezembro de 2015;60(12):3549–51.
28. Unal A, Sayharman SE, Ozel L, Unal E, Aka N, Titiz I, et al. Acute abdomen in pregnancy requiring surgical management: a 20-case series. *Eur J Obstet Gynecol Reprod Biol.* Novembro de 2011;159(1):87–90.
29. Alonso-Lázaro N, Bustamante-Balén M, Pous-Serrano S, Braithwaite-Flores A, Ponce-Romero M, Argüello-Viudez L, et al. Insertion of self-expanding metal stent for treatment of malignant obstruction in a pregnant woman. *Rev Espanola Enfermedades Dig Organo Of Soc Espanola Patol Dig.* Março de 2014;106(3):216–9.
30. Morice P, Uzan C, Uzan S. Cancer in pregnancy: a challenging conflict of interest. *The Lancet.* Fevereiro de 2012;379(9815):495–6.
31. Amant F, Halaska MJ, Fumagalli M, Dahl Steffensen K, Lok C, Van Calsteren K, et al. Gynecologic Cancers in Pregnancy: Guidelines of a Second International Consensus Meeting. *Int J Gynecol Cancer.* Março de 2014;24(3):394–403.

## Acknowledgements

I would like to acknowledge Laura Elisabete Barbosa, MD, for all the support and dedication given during this review.

# ANEXO



The Journal of Coloproctology (JCOL) publish articles that contribute to the improvement and the development of the practice, research, and training in Coloproctology and related specialities. Also published in English version, starting in vol. 31, issue 3, 2011. The guidelines are based on the format proposed by the International Committee of Medical Journal Editors (ICMJE) and published in the article: Uniform requirements for manuscripts submitted to biomedical journals, which was updated in April 2010 and is available on the Website (<http://www.icmje.org>).

## **Manuscript categories**

**Editorial** The text should have up to 900 words and 5 references.

### **Original article**

The text should have up to 3000 words, not including references and tables. It should have up to 5 tables and/or figures. The number of references should not exceed 30. Their structure should contain the following:

**Introduction:** it should be brief, defining the studied problem and highlighting its importance and gaps in knowledge.

**Methods:** the methods employed, the population studied, sources of data and selection criteria should be described in an objective and detailed manner.

Insert the protocol number of approval of the Research Ethics Committee and inform that the study was conducted according to the ethical standards required.

**Results:** they should be clearly and objectively presented, describing the obtained data only, without interpretations or comments, and, for a better understanding, they may have tables, charts and figures. The text should complement and not repeat what is described in the illustrations.

**Discussion:** it should be limited to the obtained data and results, emphasizing the new and important aspects observed in the study and discussing the agreements and disagreements with previously published studies.

**Conclusion:** it should correspond to the study objectives or assumptions, based on the results and discussion, aligned with the title, proposition and method.

### **Clinical information**

Clinical case reports, presentation of technical notes, methods and devices. They should address questions of interest to Coloproctology and related specialities.

Their structure should contain the following:

**Introduction:** it should be brief and show the theme relevance.

**Presentation of clinical case, or technique, or method, or device:** it should be described with clarity and objectiveness. It should present significant data for Coloproctology and related specialties, and have up to five figures, including tables.

**Discussion:** it should be based on the literature. The text not exceed 1500 words, not including references and figures.

Patients' initials and dates should be avoided, showing only relevant laboratorial exams for diagnosis and discussion. The total number of illustrations and/or tables should not exceed 3 and the limit of references is 20.

When the number of presented cases exceed 3, the manuscript will be classified as a Case Series, and the rules for original articles should be applicable. .

### **Review articles**

Systematic review: broad research method, conducted through a rigorous synthesis of results from original studies, either quantitative or qualitative, with the purpose of clearly answering a specific question of relevance to Coloproctology and related specialties. It should include the search strategy of original studies, the selection criteria for studies included in the review and the procedures used in the synthesis of results obtained from reviewed studies, which may or may not include meta-analysis.

Integrative review: research method that presents the synthesis of multiple published studies and enables general conclusions regarding a specific area of study, contributing to enhanced knowledge of the investigated theme. It should follow standards of methodological rigor, clarity of result presentation, enabling the reader to identify the real characteristics of studies included in the review.

Integrative review phases: elaboration of a guiding question, search strategy, data collection, critical analysis of included studies, integrative review presentation and result discussion. The text should not exceed 5000 words, not including references and tables. The total number of illustrations and tables should not exceed 8. The number of references should be limited to 60.

### **Special articles**

They should have up to 2000 words and 30 references. In all categories, in-text citation of authors should be numerical and sequential, using superscript Arabic numerals in parentheses, avoiding the indication of authors' names. In-text citations and references mentioned in legends of tables and figures should be consecutively numbered in the order of their appearance in the text, with Arabic numerals (index numbers). Only the reference number should be included, without further information.

Page charges

This journal has no page charges.

### **Submission checklist**

You can use this list to carry out a final check of your submission before you send it to the journal for review. Please check the relevant section in this Guide for Authors for more details.

### **Ensure that the following items are present:**

One author has been designated as the corresponding author with contact details:

- E-mail address
- Full postal address

All necessary files have been uploaded:

#### **Manuscript:**

- Include keywords
- All figures (include relevant captions)

- All tables (including titles, description, footnotes)
- Ensure all figure and table citations in the text match the files provided
- Indicate clearly if color should be used for any figures in print

**Graphical Abstracts / Highlights files** (where applicable)

**Supplemental files** (where applicable)

Further considerations

- Manuscript has been 'spell checked' and 'grammar checked'
- All references mentioned in the Reference List are cited in the text, and vice versa
- Permission has been obtained for use of copyrighted material from other sources (including the Internet)
- Relevant declarations of interest have been made
- Journal policies detailed in this guide have been reviewed
- Referee suggestions and contact details provided, based on journal requirements

For further information, visit our [Support Center](#).

### **Checklist ([www.jcol.org.br](http://www.jcol.org.br))**

For improved process and enhanced publication quality, we offer a checklist for your self-evaluation.



#### **Before You Begin**

### **Ethics in publishing**

Please see our information pages on [Ethics in publishing](#) and [Ethical guidelines for journal publication](#).

### **Human and animal rights**

If the work involves the use of human subjects, the author should ensure that the work described has been carried out in accordance with [The Code of Ethics of the World Medical Association](#) (Declaration of Helsinki) for experiments involving humans; [Uniform Requirements for manuscripts submitted to Biomedical journals](#). Authors should include a statement in the manuscript that informed consent was obtained for experimentation with human subjects. The privacy rights of human subjects must always be observed.

All animal experiments should comply with the [ARRIVE guidelines](#) and should be carried out in accordance with the U.K. Animals (Scientific Procedures) Act, 1986 and associated guidelines, [EU Directive 2010/63/EU for animal experiments](#), or the National Institutes of Health guide for the care and use of Laboratory animals (NIH Publications No. 8023, revised 1978) and the authors should clearly indicate in the manuscript that such guidelines have been followed.

### **Declaration of interest**

All authors must disclose any financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work. Examples of potential conflicts of interest include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding. If there are no conflicts of interest then please state this: 'Conflicts of interest: none'. [More information](#).

## Submission declaration and verification

Submission of an article implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see '[Multiple, redundant or concurrent publication](#)' section of our ethics policy for more information), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. To verify originality, your article may be checked by the originality detection service [CrossCheck](#).

## Authorship criteria

All authors should have made substantial contributions to all of the following: (1) the conception and design of the study, or acquisition of data, or analysis and interpretation of data, (2) drafting the article or revising it critically for important intellectual content, (3) final approval of the version to be submitted. Data collection and indexing are not authorship criteria. Likewise, authors are not technical assistants that perform routine tasks, physicians that refer patients or interpret routine exams and heads of services or departments not directly involved in the study. Special acknowledgments can be made to these people.

## Changes to authorship

Authors are expected to consider carefully the list and order of authors **before** submitting their manuscript and provide the definitive list of authors at the time of the original submission. Any addition, deletion or rearrangement of author names in the authorship list should be made only **before** the manuscript has been accepted and only if approved by the journal Editor. To request such a change, the Editor must receive the following from the **corresponding author**: (a) the reason for the change in author list and (b) written confirmation (e-mail, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed. Only in exceptional circumstances will the Editor consider the addition, deletion or rearrangement of authors **after** the manuscript has been accepted. While the Editor considers the request, publication of the manuscript will be suspended. If the manuscript has already been published in an online issue, any requests approved by the Editor will result in a corrigendum.

## Records of clinical essays

The **Journal of Coloproctology** supports the guideline for clinical essay recording issued by the World Health Organization (WHO) and the International Committee of Medical Journal Editors (ICMJE). Articles on clinical essays will be accepted for publication only if an identification (ID) number has been assigned by one of the Clinical Essay Record validated according to the criteria established by the WHO and ICMJE, whose addresses are at (<http://www.icmje.org>). The ID number should be displayed at the end of the abstract.

## Copyright

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' to assign to the society the copyright in the manuscript and any tables, illustrations or other material submitted for publication as part of the manuscript (the "Article") in all forms and media (whether now known or later developed), throughout the world, in all languages, for the full term of copyright, effective when the Article is accepted for publication.

### Author rights

As an author you (or your employer or institution) have certain rights to reuse your work. For more information on author rights please see <http://www.elsevier.com/copyright>.

Elsevier supports responsible sharing

Find out how you can [share your research](#) published in Elsevier journals.

Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

For non-commercial purposes, lets others distribute and copy the article, and to include in a collective work (such as an anthology), as long as they credit the author(s) and provided they do not alter or modify the article.

## Submission

Our online submission system guides you stepwise through the process of entering your article details and uploading your files. The system converts your article files to a single PDF file used in the peer-review process. Editable files (e.g., Word, LaTeX) are required to typeset your article for final publication. All correspondence, including notification of the Editor's decision and requests for revision, is sent by e-mail.

### Submit your article

Please submit your article via <http://www.evise.com/evise/jrnl/JCOL>.



### Preparation

#### Use of word processing software

It is important that the file be saved in the native format of the word processor used. The text should be in single-column format. Keep the layout of the text as

simple as possible. Most formatting codes will be removed and replaced on processing the article. In particular, do not use the word processor's options to justify text or to hyphenate words. However, do use bold face, italics, subscripts, superscripts etc. When preparing tables, if you are using a table grid, use only one grid for each individual table and not a grid for each row. If no grid is used, use tabs, not spaces, to align columns. The electronic text should be prepared in a way very similar to that of conventional manuscripts (see also the [Guide to Publishing with Elsevier](#)). Note that source files of figures, tables and text graphics will be required whether or not you embed your figures in the text. See also the section on Electronic artwork.

To avoid unnecessary errors you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your word processor.

## **Article structure**

### **The identification page**

It should contain:

a) The article title, in Portuguese and English, which should be concise and informative; it should express the manuscript content with precision. In addition, the title is important for physicians and investigators to find an article in the bibliographical databases after it is published. Please, be sure the title:

- Is not a question.
- Does not have colon or any punctuation that separates it in two parts.
- Does not reaffirm the article type. Ex.: Case Report, Review.
- Does not indicate the type of statistical analysis. Ex.: Multivariate Analysis.
- Does not include the institution name.

Full name of each author and institutional affiliation. Author affiliations should be presented in decreasing hierarchical order (e.g. Harvard University, Harvard Business School, Boston, USA) and should be written as established in its own language (e.g. Université Paris-Sorbonne; Harvard University, Universidade de São Paulo).

Name of the department and institution to which the paper should be attributed.

Name, address, e-mail of the corresponding author in charge.

Sources of support to study development.

For studies presented in scientific meetings, indicate the meeting name, place, date, type of presentation.

### **Abstract**

The second page should have the abstract, in Portuguese and English, with no more than 250 words. For original and review articles, the abstract structure should highlight the study objectives, methods, main results with significant data and conclusions. For clinical information special articles, the abstract does not need to be structured as mentioned above, but it should contain important

information for the study value recognition, as described in details in the publications: Haynes RB, Mulrow CD, Huth EJ, Altman DG, Gardiner MJ. More informative abstracts revisited. *Ann Intern Med* 1990;113:69-76 Ad Hoc Working Group for Critical Appraisal of the Medical Literature. A proposal for more informative abstracts of clinical articles. *Ann Intern Med* 1987;106:598-604.

## **Keywords**

After the abstract, specify three to six terms in Portuguese and in English the subject of the study should be included as well as the corresponding. Keywords in must be based on the Health and Science Keywords (DeCS), published by Bireme and available at (<http://decs.bvs.br>), and Medical Subject Headings (MeSH) is the Nation Library Medicine controlled vocabulary thesaurus used for indexing articles for PubMed at (<http://www.nlm.nih.gov/mesh/meshhome.html>).

Abbreviations should be indicated when they first appear in the text. After that, the full name should not be repeated.

## **Acknowledgements**

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

## **Statistical analysis**

The authors should demonstrate that the statistical procedures used in the study were not only appropriate to test the study hypotheses, but also correctly interpreted. The levels of statistical significance (ex.  $p < 0.05$ ;  $p < 0.01$ ;  $p < 0.001$ ) should be mentioned.

## **Electronic artwork**

### **General points**

- Make sure you use uniform lettering and sizing of your original artwork.
- Embed the used fonts if the application provides that option.
- Aim to use the following fonts in your illustrations: Arial, Courier, Times New Roman, Symbol, or use fonts that look similar.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Provide captions to illustrations separately.
- Size the illustrations close to the desired dimensions of the published version.
- Submit each illustration as a separate file.

A detailed [guide on electronic artwork](#) is available.

**You are urged to visit this site; some excerpts from the detailed information are given here.**

### **Formats**

If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is' in the native document format.



Regardless of the application used other than Microsoft Office, when your electronic artwork is finalized, please 'Save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings, embed all used fonts.

TIFF (or JPEG): Color or grayscale photographs (halftones), keep to a minimum of 300 dpi.

TIFF (or JPEG): Bitmapped (pure black & white pixels) line drawings, keep to a minimum of 1000 dpi.

TIFF (or JPEG): Combinations bitmapped line/half-tone (color or grayscale), keep to a minimum of 500 dpi.

**Please do not:**

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); these typically have a low number of pixels and limited set of colors;
- Supply files that are too low in resolution;
- Submit graphics that are disproportionately large for the content.

**Figure captions**

Ensure that each illustration has a caption. Supply captions separately, not attached to the figure. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

**Figures**

The illustrations (pictures, charts, drawings, etc.) should be submitted individually. They should be consecutively numbered, with Arabic numerals, in the order of their appearance in the text, and they should be clear enough to enable their reproduction. Photocopies will not be accepted.

**Tables**

Please submit tables as editable text and not as images. Tables can be placed either next to the relevant text in the article, or on separate page(s) at the end. Number tables consecutively in accordance with their appearance in the text and place any table notes below the table body. Be sparing in the use of tables and ensure that the data presented in them do not duplicate results described elsewhere in the article. Please avoid using vertical rules and shading in table cells.

**References**

**Citation in text**

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication.



## Reference style

They should be consecutively numbered in the order of their appearance in the text and identified with Arabic numerals. They should be presented according to the style presented by the List of Journal Indexed Medicus, of the National Library of Medicine, which can be accessed at <http://www.nlm.gov/tsd/serials/lji.html>. The authors should be sure that in-text citations of references are included in the list of references with exact dates and correctly spelled names of authors. The accuracy of references is the authors' responsibility. Personal notes, unprecedented studies or studies in progress may be cited when really required, but should not be included in the list of references; only cited in the text or footnotes. Cite up to six authors for each reference. If any reference has more than six authors, cite the six first names, followed by "et al.". We request texts with lean writing style. Shorter texts involve shorter revision and formatting times, and have higher chances of quick publication.



## After Acceptance

### Online proof correction

Corresponding authors will receive an e-mail with a link to our online proofing system, allowing annotation and correction of proofs online. The environment is similar to MS Word: in addition to editing text, you can also comment on figures/tables and answer questions from the Copy Editor. Web-based proofing provides a faster and less error-prone process by allowing you to directly type your corrections, eliminating the potential introduction of errors.

If preferred, you can still choose to annotate and upload your edits on the PDF version. All instructions for proofing will be given in the e-mail we send to authors, including alternative methods to the online version and PDF.

We will do everything possible to get your article published quickly and accurately. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. It is important to ensure that all corrections are sent back to us in one communication. Please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.



## Author Inquiries

Visit the [Elsevier Support Center](#) to find the answers you need. Here you will find everything from Frequently Asked Questions to ways to get in touch. You can also [check the status of your submitted article](#) or find out [when your accepted article will be published](#).